

## Section 2: Simultaneous Equations

### Solutions to Exercise

1. (i)  $2x + 5y = 11$  (1)

$2x - y = 5$  (2)

Subtracting:  $6y = 6$

$y = 1$

Substituting into (1):  $2x + 5 \times 1 = 11$

$2x = 6$

$x = 3$

The solution is  $x = 3, y = 1$ . Check:  $2x + 5y = 2 \times 3 + 5 \times 1 = 11$

$2x - y = 2 \times 3 - 1 = 5$

(ii)  $x + 2y = 6$  (1)  $\times 4$   $4x + 8y = 24$

$4x + 3y = 4$  (2)  $4x + 3y = 4$

Subtracting:  $5y = 20$

$y = 4$

Substituting into (1):  $x + 2 \times 4 = 6$

$x = -2$

The solution is  $x = -2, y = 4$ . Check:  $x + 2y = -2 + 8 = 6$

$4x + 3y = -8 + 12 = 4$

(iii)  $3a - 2b = 4$  (1)  $\times 2$   $6a - 4b = 8$

$5a + 4b = 3$  (2)  $5a + 4b = 3$

Adding:  $11a = 11$

$a = 1$

Substituting into (1):  $3 \times 1 - 2b = 4$

$-2b = 1$

$b = -\frac{1}{2}$

The solution is  $a = 1, b = -\frac{1}{2}$ . Check:  $3a - 2b = 3 + 1 = 4$

$5a + 4b = 5 - 2 = 3$

(iv)  $2p - 5q = 5$  (1)  $\times 3$   $6p - 15q = 15$

$3p - 2q = -9$  (2)  $\times 2$   $6p - 4q = -18$

Subtracting:  $-11q = 33$

$q = -3$

Substituting into (1):  $2p - 5 \times -3 = 5$

$$2p = -10$$

$$p = -5$$

The solution is  $p = -5$ ,  $q = -3$ . Check:  $2p - 5q = -10 + 15 = 5$

$$3p - 2q = -15 + 6 = -9$$

(v)  $5x + 3y = 9$  (1)

$y = 3x - 4$  (2)

Substituting (2) into (1):  $5x + 3(3x - 4) = 9$

$$5x + 9x - 12 = 9$$

$$14x = 21$$

$$x = \frac{3}{2}$$

Substituting into (1):  $y = 3 \times \frac{3}{2} - 4 = \frac{9}{2} - 4 = \frac{1}{2}$

The solution is  $x = \frac{3}{2}$ ,  $y = \frac{1}{2}$ . Check:  $5x + 3y = \frac{15}{2} + \frac{3}{2} = 9$

(vi)  $3a + 2b = 1$  (1)  $\times 2$   $6a + 4b = 2$

$9a - 4b = 4$  (2)  $9a - 4b = 4$

Adding:  $15a = 6$

$$a = \frac{2}{5}$$

Substituting into (1):  $3 \times \frac{2}{5} + 2b = 1$

$$2b = 1 - \frac{6}{5} = -\frac{1}{5}$$

$$b = -\frac{1}{10}$$

The solution is  $a = \frac{2}{5}$ ,  $b = -\frac{1}{10}$ . Check:  $3a + 2b = \frac{6}{5} - \frac{1}{5} = 1$

$$9a - 4b = \frac{18}{5} + \frac{2}{5} = 4$$

2. (i)  $7x^2 + y^2 = 64$  (1)

$x + y = 4$  (2)

(2)  $\Rightarrow y = 4 - x$

Substituting into (1):  $7x^2 + (4 - x)^2 = 64$

$$7x^2 + 16 - 8x + x^2 = 64$$

$$8x^2 - 8x - 48 = 0$$

$$x^2 - x - 6 = 0$$

$$(x - 3)(x + 2) = 0$$

$$x = 3 \text{ or } x = -2$$

When  $x = 3$ ,  $y = 4 - 3 = 1$

When  $x = -2$ ,  $y = 4 - (-2) = 6$

The solutions are  $x = 3$ ,  $y = 1$  and  $x = -2$ ,  $y = 6$

Check:  $x = 3, y = 1 \Rightarrow 7x^2 + y^2 = 63 + 1 = 64$

$x = -2, y = 6 \Rightarrow 7x^2 + y^2 = 28 + 36 = 64$

(ii)  $3x^2 - 2y^2 = -5$  (1)

$y - x = 1$  (2)

(2)  $\Rightarrow y = 1 + x$

Substituting into (1):  $3x^2 - 2(1 + x)^2 = -5$

$$3x^2 - 2(1 + 2x + x^2) = -5$$

$$3x^2 - 2 - 4x - 2x^2 = -5$$

$$x^2 - 4x + 3 = 0$$

$$(x - 1)(x - 3) = 0$$

$$x = 1 \text{ or } x = 3$$

When  $x = 1$ ,  $y = 1 + 1 = 2$

When  $x = 3$ ,  $y = 1 + 3 = 4$

The solutions are  $x = 1$ ,  $y = 2$  and  $x = 3$ ,  $y = 4$

Check:  $x = 1, y = 2 \Rightarrow 3x^2 - 2y^2 = 3 - 5 = -5$

$x = 3, y = 4 \Rightarrow 3x^2 - 2y^2 = 27 - 32 = -5$

(iii)  $p^2 + pq = 2$  (1)

$q - p = 3$  (2)

(2)  $\Rightarrow q = 3 + p$

Substituting into (1):  $p^2 + p(3 + p) = 2$

$$p^2 + 3p + p^2 = 2$$

$$2p^2 + 3p - 2 = 0$$

$$(2p - 1)(p + 2) = 0$$

$$p = \frac{1}{2} \text{ or } p = -2$$

When  $p = \frac{1}{2}$ ,  $q = 3 + \frac{1}{2} = \frac{7}{2}$

When  $p = -2$ ,  $q = 3 - 2 = 1$

The solutions are  $p = \frac{1}{2}$ ,  $q = \frac{7}{2}$  and  $p = -2$ ,  $q = 1$ .

Check:  $p = \frac{1}{2}, q = \frac{7}{2} \Rightarrow p^2 + pq = \frac{1}{4} + \frac{7}{4} = 2$

$p = -2, q = 1 \Rightarrow p^2 + pq = 4 - 2 = 2$

(iv)  $8a^2 - b^2 = 2$  (1)

$2a + b = 1$  (2)

(2)  $\Rightarrow b = 1 - 2a$

Substituting into (1):  $8a^2 - (1 - 2a)^2 = 2$

$$8a^2 - (1 - 4a + 4a^2) = 2$$

$$8a^2 - 1 + 4a - 4a^2 = 2$$

$$4a^2 + 4a - 3 = 0$$

$$(2a + 3)(2a - 1) = 0$$

$$a = -\frac{3}{2} \text{ or } a = \frac{1}{2}$$

When  $a = -\frac{3}{2}$ ,  $b = 1 - 2 \times -\frac{3}{2} = 1 + 3 = 4$

When  $a = \frac{1}{2}$ ,  $b = 1 - 2 \times \frac{1}{2} = 1 - 1 = 0$

The solutions are  $a = -\frac{3}{2}$ ,  $b = 4$  and  $a = \frac{1}{2}$ ,  $b = 0$ .

Check:  $a = -\frac{3}{2}$ ,  $b = 4 \Rightarrow 8a^2 - b^2 = 8 \times \frac{9}{4} - 16 = 18 - 16 = 2$

$a = \frac{1}{2}$ ,  $b = 0 \Rightarrow 8a^2 - b^2 = 8 \times \frac{1}{4} - 0 = 2$

3.  $xy - 9 = 15$

$2x + 2y = 20$

$y = 10 - x$

$x(10 - x) = 24$

$10x - x^2 = 24$

$x^2 - 10x + 24 = 0$

$(x - 6)(x - 4) = 0$

$x = 6$  and  $y = 4$  (or  $x = 4$  and  $y = 6$ )

4. (i)

(1)  $x - y + z = 4$

(2)  $3x + y + 3z = -4$

(3)  $x + y + 2z = -2$

eliminate  $x$  by combining equations

(3) - (1)  $2y + z = -6$

3 \* (3) - (2)  $2y + 3z = -2$

eliminate  $y$  by combining equations

$2z = -2 - (-6)$

$z = 2$

Substitute in (1) and (3)

$$(1) \quad x - y + 2 = 4$$

$$(3) \quad x + y + 4 = -2$$

$$2x + 6 = 2$$

$$2x = -4$$

$$x = -2$$

Substitute in (3)

$$(3) \quad -2 + y + 4 = -2$$

$$y = -4$$

so  $x = -2$ ,  $y = -4$  and  $z = 2$

(ii)

$$(1) \quad 2x + 2y - z = 0$$

$$(2) \quad 8x - 12y - z = -32$$

$$(3) \quad -2x - 4y - z = 6$$

eliminate  $z$  by combining equations

$$(1) - (3) \quad 4x + 6y = -6$$

$$(2) - (1) \quad 6x - 14y = -32$$

eliminate  $x$  by combining equations

$$6x + 9y = -9$$

$$6x - 14y = -32$$

$$9y + 14y = -9 - (-32)$$

$$23y = 23$$

$$y = 1$$

Substitute in (1) and (3)

$$(1) \quad 2x + 2 - z = 0$$

$$(3) \quad -2x - 4 - z = 6$$

$$-2 - 2z = 6$$

$$2z = -8$$

$$z = -4$$

Substitute in (1)

$$2x + 2 + 4 = 0$$

$$2x = -6$$

$$x = -3$$

so  $x = -3$ ,  $y = 1$  and  $z = -4$

5.

(i)

$$(1) \quad 2x + y + 3z = 8$$

$$(2) \quad -3x + 3y + 2z = 12$$

$$(3) \quad 3x + 2y - z = -4$$

eliminate  $y$  by combining equations

$$3 \times (1) - (2) \quad 9x + 7z = 12$$

$$2 \times (1) - (3) \quad x + 7z = 20$$

eliminate  $z$  by combining equations

$$8x = -8$$

$$x = -1$$

Substitute in (1) and (2) and combine

$$(1) \quad -2 + y + 3z = 8$$

$$(2) \quad 3 + 3y + 2z = 12$$

$$7z = 21$$

$$z = 3$$

Substitute in (3)

$$(1) \quad -2 + y + 9 = 8$$

$$y = 1$$

so  $x = -1$ ,  $y = 1$  and  $z = 3$

(ii)

$$(1) \quad x - 2y - 2z = 1$$

$$(2) \quad 2x - y - 14z = 3$$

$$(3) \quad 2x - 3y + 3z = -8$$

eliminate  $x$  by combining equations

$$(2) - (3) \quad 2y - 17z = 11$$

$$2 \times (1) - (3) \quad -y - 7z = 10$$

eliminate  $y$  by combining equations

$$-31z = 31$$

$$z = -1$$

Substitute in (1) and (2) and combine

$$(1) \quad x - 2y + 2 = 1$$

$$(2) \quad 2x - y + 14 = 3$$

$$-3y = 9$$

$$y = -3$$

Substitute in (3)

$$(1) \quad x + 6 + 2 = 1$$

$$x = -7$$

so  $x = -7$ ,  $y = -3$  and  $z = -1$

6. let  $x$  represent number of cupcakes,  $y$  be number of bags crisps and  $z$  be number of cans of soda

Abby:  $16x + 40y + 24z = 36$

Ben:  $24x + 20y + 12z = 30$

Carrie:  $12x + 10y + 36z = 30$

Simplify the equations:

(A):  $4x + 10y + 6z = 9$

(B):  $12x + 10y + 6z = 15$

(C):  $6x + 5y + 18z = 15$

eliminate  $y$  and  $z$  by combining equations

(B) - (A)  $8x = 6$

$$x = \frac{6}{8} = \frac{3}{4} = 0.75$$

Substitute in (A) and (C) and combine

(A)  $4(0.75) + 10y + 6z = 9 \rightarrow (A) \quad 10y + 6z = 6$

(C)  $6(0.75) + 5y + 18z = 15 \rightarrow (C) \quad 5y + 18z = 10.5$

$$30z = 15$$

$$z = \frac{15}{30} = 0.5$$

Substitute in (A)

(1)  $10y + 3 = 6$

$$10y = 3$$

$$y = \frac{3}{10} = 0.3$$

So the cupcakes were 75p each, the crisps were 30p a bag and the sodas were 50p a can.